

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Sheep anti Guinea Pig IgM (Fc specific), Clone: [Polyclonal], Monoclonal NMB-SHAGP/IGM(FC)

| | |
|--------------------------|--|
| Artikelname | Sheep anti Guinea Pig IgM (Fc specific), Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-SHAGP/IGM(FC) |
| Hersteller Artikelnummer | ShAGp/IgM(Fc) |
| Alternativnummer | NMB-SHAGP/IGM(FC) |
| Hersteller | NordicMubio |
| Wirt | Sheep |
| Kategorie | Antikörper |
| Produktbeschreibung | The reactivity of the antiserum is restricted to the Fc part of the IgM molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal guinea pig serum a single precipitin line is obtained which s... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| Puffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added. Reconstitute the lyophilized a |
| Quelle | Highly purified normal IgM isolated from pooled guinea pig serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|------------------------|--|
| Formel | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added. |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl in a 3 mm diameter center well and 2 µl serum samples (neat and serially d |