

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Swine anti Goat IgM (Fc specific), Clone: [Polyclonal], Porcine, Monoclonal NMB-SWAG/IGM(FC)

Artikelname	Swine anti Goat IgM (Fc specific), Clone: [Polyclonal], Porcine, Monoclonal
Artikelnummer	NMB-SWAG/IGM(FC)
Hersteller Artikelnummer	SwAG/IgM(Fc)
Alternativnummer	NMB-SWAG/IGM(FC)
Hersteller	NordicMubio
Wirt	Porcine
Kategorie	Antikörper
Produktbeschreibung	The reactivity of the antiserum is restricted to the Fc part of the IgM molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal goat serum a single precipitin line is obtained which shows a...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]
Puffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added. Reconstitute the lyophilized
Quelle	Highly purified normal IgM isolated from pooled goat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

Formel	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added.
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl serum, plasma or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samp