

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Swine anti Human C3c, Clone: [Polyclonal], Porcine, Monoclonal NMB-SWAHU/C3C

| | |
|--------------------------|--|
| Artikelname | Swine anti Human C3c, Clone: [Polyclonal], Porcine, Monoclonal |
| Artikelnummer | NMB-SWAHU/C3C |
| Hersteller Artikelnummer | SwAHu/C3c |
| Alternativnummer | NMB-SWAHU/C3C |
| Hersteller | NordicMubio |
| Wirt | Porcine |
| Kategorie | Antikörper |
| Produktbeschreibung | In immunoelectrophoresis against fresh human serum, a single precipitin line is obtained in the beta-1 region representing native C3. Against serum containing partly activated C3, a precipitin line is obtained which extends from the beta-1 into the a... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| NCBI | 718 |
| UniProt | P01024 |
| Puffer | Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of normal pooled swine serum. No foreign proteins added. Reconstitute the lyophilized anti |

| | |
|------------------------|--|
| Quelle | C3c is the major fragment resulting from the C3 cleavage by C3 convertase and factor i. It is composed of an intact beta chain bound to two fragments of the alpha chain. The protein is isolated and purified from pooled normal human serum by precipitation |
| Formel | Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of normal pooled swine serum. No foreign proteins added. |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 µl human plasma or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl plasma samples (neat and se |