

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-p53 Purified, Clone: [BP53-12], Monoclonal EXB-11-114-C100

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Anti-p53 Purified, Clone: [BP53-12], Monoclonal |
| Artikelnummer | EXB-11-114-C100 |
| Hersteller Artikelnummer | 11-114-C100 |
| Alternativnummer | EXB-11-114-C100 |
| Hersteller | EXBIO |
| Kategorie | Antikörper |
| Applikation | ELISA, FC, ICC, IHC-P, IP, WB |
| Spezies Reaktivität | Human, Primate |
| Immunogen | Bacterially expressed full-length wild-type p53 |
| Produktbeschreibung | The tumour suppressor protein p53 is a key element of intracellular anticancer protection. It mediates cell cycle arrest or apoptosis in response to DNA damage or to starvation for pyrimidine nukleotides. It is up-regulated in response to these stres... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/ml |
| Klon-Bezeichnung | [BP53-12] |
| Isotyp | Mouse IgG2a |
| Puffer | Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Lagerung | 2°C to 8°C |

| | |
|------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target-Kategorie | p53 |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western blotting: Recommended dilution: 1-2 µg/ml, overnight at 4C, positive control: RAMOS human lymphoma cell line, non-reducing conditions. SDS-PAGE (12% separating gel). Immunohistochemistry: Recommended dilution: 5-10 µg/ml. Flow cytometry: |