

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Hu CD79a PE, Clone: [HM47], Monoclonal EXB-1P-731-T100

|                          |   |
|--------------------------|---|
| Artikelname              | Anti-Hu CD79a PE, Clone: [HM47], Monoclonal   |
| Artikelnummer            | EXB-1P-731-T100   |
| Hersteller Artikelnummer | 1P-731-T100   |
| Alternativnummer         | EXB-1P-731-T100   |
| Hersteller               | EXBIO   |
| Kategorie                | Antikörper  |
| Applikation              | FC  |
| Spezies Reaktivität      | Bovine, Canine, Equine, Gallus, Guinea pig, Human, Mouse, Porcine, Primate, Rabbit, Rat   |
| Immunogen                | Synthetic peptide corresponding to C terminal amino acids 208-222 of human CD79a  |
| Konjugation              | PE  |
| Produktbeschreibung      | CD79a (Ig alpha, MB1) forms disulfide-linked heterodimer with CD79b (Ig beta). They both are transmembrane proteins with extended cytoplasmic domains containing immunoreceptor tyrosine activation motives (ITAMs), and together with cell surface immuno... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [HM47]  |
| Isotyp                   | Mouse IgG1 kappa  |

|                        |  |
|------------------------|--|
| Puffer                 | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  |
| Lagerung               | 2°C to 8°C   |
| Target-Kategorie       | CD79a  |
| Antibody Type          | Monoclonal Antibody  |
| Application Verdünnung | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.                            |
| Anwendungsbeschreibung | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.<br>Intracellular staining. |