

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Hu CD41a Pacific Blue(TM), Clone: [HIP8], Monoclonal EXB-PB-417-T100

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Anti-Hu CD41a Pacific Blue(TM), Clone: [HIP8], Monoclonal |
| Artikelnummer | EXB-PB-417-T100 |
| Hersteller Artikelnummer | PB-417-T100 |
| Alternativnummer | EXB-PB-417-T100 |
| Hersteller | EXBIO |
| Kategorie | Antikörper |
| Applikation | FC |
| Spezies Reaktivität | Human, Primate |
| Konjugation | Pacific Blue |
| Produktbeschreibung | CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a v... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [HIP8] |
| Isotyp | Mouse IgG1 |
| Puffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Lagerung | 2°C to 8°C |

| | |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target-Kategorie | CD41a |
| Antibody Type | Monoclonal Antibody |
| Application Verdünnung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Anwendungsbeschreibung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |