

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Hu Ig Kappa Light Chain Pacific Blue(TM), Clone: [A8B5], Monoclonal EXB-PB-504-T100

|                          |   |
|--------------------------|---|
| Artikelname              | Anti-Hu Ig Kappa Light Chain Pacific Blue(TM), Clone: [A8B5], Monoclonal  |
| Artikelnummer            | EXB-PB-504-T100   |
| Hersteller Artikelnummer | PB-504-T100   |
| Alternativnummer         | EXB-PB-504-T100   |
| Hersteller               | EXBIO   |
| Kategorie                | Antikörper  |
| Applikation              | FC  |
| Spezies Reaktivität      | Human   |
| Konjugation              | Pacific Blue  |
| Produktbeschreibung      | Immunoglobulin classes share the same basic four polypeptide chain structure of two heavy chains (five heavy chains types) and two light chains (kappa, lambda, both having a molecular weight of 22.5kDa). Kappa and lambda consist of a variable region ... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [A8B5]  |
| Isotyp                   | Mouse IgG1  |
| Puffer                   | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide   |
| Lagerung                 | 2°C to 8°C  |

|                        |  |
|------------------------|--|
| Target-Kategorie       | Ig Kappa Light Chain   |
| Antibody Type          | Monoclonal Antibody  |
| Application Verdünnung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Anwendungsbeschreibung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |