

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-HLA-DR APC-Cy(TM)7, Clone: [L243], APC/Cy7, Monoclonal EXB-T4-690-T100

| | |
|--------------------------|---|
| Artikelname | Anti-HLA-DR APC-Cy(TM)7, Clone: [L243], APC/Cy7, Monoclonal |
| Artikelnummer | EXB-T4-690-T100 |
| Hersteller Artikelnummer | T4-690-T100 |
| Alternativnummer | EXB-T4-690-T100 |
| Hersteller | EXBIO |
| Kategorie | Antikörper |
| Applikation | FC |
| Spezies Reaktivität | Canine, Human, Primate |
| Immunogen | Human B lymphocytes |
| Konjugation | APC/Cy7 |
| Produktbeschreibung | HLA-DR, a member of MHC class II glycoproteins, that bind intracellularly processed peptides and present them to the Th cells, is composed of 36 kDa alpha chain and 27 kDa beta chain, both anchored in the plasma membrane. Together with other MHC II m... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [L243] |
| Isotyp | Mouse IgG2a kappa |
| Puffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |

| | |
|------------------------|--|
| Lagerung | 2°C to 8°C |
| Target-Kategorie | HLA-DR |
| Antibody Type | Monoclonal Antibody |
| Application Verdünnung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Anwendungsbeschreibung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |