

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Hu CD27 PE-Cy(TM)5, Clone: [LT27], PE/Cy5, Monoclonal EXB-T8-308-T100

|                          |   |
|--------------------------|---|
| Artikelname              | Anti-Hu CD27 PE-Cy(TM)5, Clone: [LT27], PE/Cy5, Monoclonal  |
| Artikelnummer            | EXB-T8-308-T100   |
| Hersteller Artikelnummer | T8-308-T100   |
| Alternativnummer         | EXB-T8-308-T100   |
| Hersteller               | EXBIO   |
| Kategorie                | Antikörper  |
| Applikation              | FC  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human peripheral blood lymphocytes  |
| Konjugation              | PE/Cy5  |
| Produktbeschreibung      | CD27 is a transmembrane 55 kDa protein of the nerve growth factor-receptor family, expressed as a disulfide-linked homodimer on mature thymocytes, peripheral blood T cells and a subpopulation of B cells. Activation of T cells via TCR-CD3 complex resu... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [LT27]  |
| Isotyp                   | Mouse IgG2a   |
| Puffer                   | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide   |

|                        |  |
|------------------------|--|
| Lagerung               | 2°C to 8°C   |
| Target-Kategorie       | CD27   |
| Antibody Type          | Monoclonal Antibody  |
| Application Verdünnung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Anwendungsbeschreibung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |