

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Myosin (Skeletal, Slow) Monoclonal Antibody (Clone: NOQ7.5.4D), Clone: [NOQ7.5.4D], Mouse ABI-39-1070

| | |
|--------------------------|---|
| Artikelname | Anti-Myosin (Skeletal, Slow) Monoclonal Antibody (Clone: NOQ7.5.4D), Clone: [NOQ7.5.4D], Mouse |
| Artikelnummer | ABI-39-1070 |
| Hersteller Artikelnummer | 39-1070 |
| Alternativnummer | ABI-39-1070-100UG,ABI-39-1070-100UG/VIAL |
| Hersteller | Abeomics |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | WB |
| Spezies Reaktivität | Human |
| Immunogen | Human skeletal muscle myosin purified from myofibrils. |
| Produktbeschreibung | Myosin is composed of 2 heavy chains of about 200,000 daltons each and 4 light chains of about 20,000 daltons each. Skeletal Myosin(slow),, also known as light chain 3(MYL3), mapped to 3p. Fodor et al.(1989) found that the MYL3 gene has 7 exons, the ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [NOQ7.5.4D] |
| Isotyp | Mouse IgG1 |
| NCBI | 4625 |

| | |
|------------------------|--|
| UniProt | P12883 |
| Puffer | Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN ₃ as preservative. Reconstitute : Add 1ml of PBS buffer will yield a concentration of 100 µg/ml. |
| Reinheit | Ascites |
| Formel | Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN ₃ as preservative. Reconstitute : Add 1ml of PBS buffer will yield a concentration of 100ug/ml. |
| Target-Kategorie | Myosin-7 |
| Application Verdünnung | Western blot : 0.5-2µg/ml, Immunohistochemistry(Paraffin-embedded Section) : 1-2µg/ml |