

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Tau Monoclonal Antibody (Clone: TAU-2), Clone: [TAU-2], Mouse ABI-39-1093

|                          |   |
|--------------------------|---|
| Artikelname              | Anti-Tau Monoclonal Antibody (Clone: TAU-2), Clone: [TAU-2], Mouse  |
| Artikelnummer            | ABI-39-1093   |
| Hersteller Artikelnummer | 39-1093   |
| Alternativnummer         | ABI-39-1093-100UG,ABI-39-1093-100UG/VIAL  |
| Hersteller               | Abeomics  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC-P, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Bovine microtubule-associated proteins(MAPs).   |
| Produktbeschreibung      | The microtubule-associated proteins (MAPs) coassemble with tubulin into microtubules in vitro. Microtubule-associated protein tau appears to be enriched in axons. Tau are composed of 352 to 441 amino acids. The isoforms differ from each other by the ... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [TAU-2]   |
| Isotyp                   | Mouse IgG1  |
| NCBI                     | <a href="#">281296</a>  |
| UniProt                  | <a href="#">P29172</a>  |

|                        |  |
|------------------------|--|
| Puffer                 | Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN3 as preservative. Reconstitute : Add 1ml of PBS buffer will yield a concentration of 100 µg/ml. |
| Reinheit               | Ascites  |
| Formel                 | Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN3 as preservative. Reconstitute : Add 1ml of PBS buffer will yield a concentration of 100ug/ml.  |
| Target-Kategorie       | Microtubule-associated protein tau   |
| Application Verdünnung | Western blot : 0.5-1µg/ml, Immunohistochemistry(Paraffin-embedded Section) : 1-2µg/ml  |