

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant Mouse Monoclonal Antibody to Ep-CAM / CD326 (Epithelial Marker)(Clone : rMOC-31), Clone: [rMOC-31] ABI-12-1006**

Artikelname	Recombinant Mouse Monoclonal Antibody to Ep-CAM / CD326 (Epithelial Marker)(Clone : rMOC-31), Clone: [rMOC-31]
Artikelnummer	ABI-12-1006
Hersteller Artikelnummer	12-1006
Alternativnummer	ABI-12-1006-100UG
Hersteller	Abeomics
Kategorie	Antikörper
Applikation	FACS, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Neuraminidase treated GLS-1 human small cell lung carcinoma cells
Produktbeschreibung	Binding epitope of this antibody is located in the first EGF-like repeat domain (EGF1) between amino acids 27-59 of Ep-CAM. EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial...
Klonalität	Monoclonal
Klon-Bezeichnung	[rMOC-31]
Isotyp	Mouse IgG1, kappa
NCBI	<a href="#">4072</a>
UniProt	<a href="#">P16422</a>

Puffer	200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Reinheit	Purified Ab with BSA and Azide at 200ug/ml
Formulierung	Purified
Formel	200 µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Target-Kategorie	Ep-CAM / CD326 (Epithelial Marker)
Application Verdünnung	Flow Cytometry (1-2µg/million cells), Immunofluorescence (1-2µg/ml), Western Blot (1-2µg/ml), Immunohistochemistry (Formalin-fixed) (1-2µg/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1m