

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Hu CD271 APC-Cy(TM)7, Clone: [NGFR5], APC/Cy7, Monoclonal EXB-T4-642-T100

| | |
|--------------------------|---|
| Artikelname | Anti-Hu CD271 APC-Cy(TM)7, Clone: [NGFR5], APC/Cy7, Monoclonal |
| Artikelnummer | EXB-T4-642-T100 |
| Hersteller Artikelnummer | T4-642-T100 |
| Alternativnummer | EXB-T4-642-T100 |
| Hersteller | EXBIO |
| Kategorie | Antikörper |
| Applikation | FC |
| Spezies Reaktivität | Feline, Ferret, Human, Primate, Rabbit |
| Immunogen | Purified CD271 protein isolated from human melanoma cell line A875 |
| Konjugation | APC/Cy7 |
| Produktbeschreibung | CD271 / NGFR, also known as p75NGFR or p75NTR, is a 75 kDa low affinity receptor for the NGF (nerve growth factor), BDNF (brain-derived growth factor), and other neurotrophins, such as NT3 and NT4/5. Unlike other members of the tumor necrosis factor ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [NGFR5] |
| Isotyp | Mouse IgG1 |
| Puffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |

| | |
|------------------------|--|
| Target-Kategorie | CD271 |
| Antibody Type | Monoclonal Antibody |
| Application Verdünnung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Anwendungsbeschreibung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. Extracellular and intracellular stai |