

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti-Rabbit IgG (H&L), TRITC conjugated, Polyclonal AGR-AS10-1196

| | |
|--------------------------|--|
| Artikelname | Goat anti-Rabbit IgG (H&L), TRITC conjugated, Polyclonal |
| Artikelnummer | AGR-AS10-1196 |
| Hersteller Artikelnummer | AS10-1196 |
| Alternativnummer | AGR-AS10-1196 |
| Hersteller | Agrisera |
| Wirt | Goat |
| Kategorie | Antikörper |
| Konjugation | tritic |
| Produktbeschreibung | Goat anti-rabbit IgG (H&L), Rhodamine (TRITC-tetramethyl rhodamine-5-isothiocyanate) conjugated is a secondary antibody which binds to rabbit IgGs in immunological assays. TRITC has $\lambda_{max} = 550 \text{ nm}$, $\lambda_{em} = 570 \text{ nm}$. Antibodies are affinity purified using... |
| Klonalität | Polyclonal |
| Reinheit | Immunogen affinity purified IgG. |
| Formulierung | Lyophilized |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | The optimal working dilution should be determined by the investigator, (ICC), immunohistochemistry (IHC) (Flow cyt) |

Anwendungsbeschreibung

For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily