

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat Antiserum anti-Bovine IgG (F(ab)2)-unconj., MinX none, Polyclonal , Unconjugated DNA-SEC-182408

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Goat Antiserum anti-Bovine IgG (F(ab)2)-unconj., MinX none, Polyclonal , Unconjugated |
| Artikelnummer | DNA-SEC-182408 |
| Hersteller Artikelnummer | SEC-182408 |
| Alternativnummer | DNA-SEC-182408 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | WB, IHC, ELISA |
| Spezies Reaktivität | Bovine |
| Immunogen | Bovine IgG F(ab)2 fragment |
| Konjugation | Unconjugated |
| Produktbeschreibung | Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglu... |
| Klonalität | Polyclonal |
| Konzentration | 100 mg/mL |
| Isotyp | Ig |

| | |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-goat serum, Bovine IgG, Bovine IgG F(ab)2 and Bovine Serum. No reaction was observed |
| Formel | 20 mM K3PO4, 150 mM NaCl, pH 7.2, lyophilisate, 0.01% NaN3 |
| Target-Kategorie | Bovine |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | ELISA Dilution: 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Anwendungsbeschreibung | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maxi |