

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Donkey Antiserum anti-Rabbit IgG (Fc)-unconj., MinX none, Polyclonal DNA-SEC-182459

| | |
|--------------------------|---|
| Artikelname | Donkey Antiserum anti-Rabbit IgG (Fc)-unconj., MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-182459 |
| Hersteller Artikelnummer | SEC-182459 |
| Alternativnummer | DNA-SEC-182459 |
| Hersteller | dianova |
| Wirt | Donkey |
| Kategorie | Antikörper |
| Applikation | WB, IHC, ELISA |
| Spezies Reaktivität | Rabbit |
| Immunogen | Rabbit IgG F(c) fragment |
| Konjugation | Unconjugated |
| Produktbeschreibung | Anti-Rabbit IgG F(c) generated in donkey is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of rabbit IgG and... |
| Klonalität | Polyclonal |
| Konzentration | 80 mg/mL |
| Isotyp | Ig |

| | |
|------------------------|--|
| Puffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-donkey serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit Serum. No reaction was observe |
| Formel | 10 mM NaPO ₄ ,150 mM NaCl,pH 7,2,lyophilisate,Azide/BSA free |
| Target-Kategorie | Rabbit |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:2,000 - 1:10,000 |
| Anwendungsbeschreibung | Antibody Anti-Rabbit IgG F(c) is suitable for immunoblotting (western or dot blot), ELISA, and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency. |