

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat Antiserum anti-Rat IgG (H+L)-unconj., MinX none, Polyclonal DNA-SEC-182460

|                          |   |
|--------------------------|---|
| Artikelname              | Goat Antiserum anti-Rat IgG (H+L)-unconj., MinX none, Polyclonal  |
| Artikelnummer            | DNA-SEC-182460  |
| Hersteller Artikelnummer | SEC-182460  |
| Alternativnummer         | DNA-SEC-182460  |
| Hersteller               | dianova   |
| Wirt                     | Goat  |
| Kategorie                | Antikörper  |
| Applikation              | WB, IHC, ELISA  |
| Spezies Reaktivität      | Rat   |
| Immunogen                | Rat IgG whole molecule  |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | Anti-Rat IgG (H&L) generated in goat detects rat Immunoglobulin G. Both the Heavy and Light chains of the antibody molecule are present. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the c... |
| Klonalität               | Polyclonal  |
| Konzentration            | unprocessed   |
| Isotyp                   | Ig  |

|                        |   |
|------------------------|---|
| Puffer                 | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |
| Reinheit               | This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-goat serum, Rat IgG and Rat Serum. |
| Formel                 | 10 mM NaPO <sub>4</sub> , 150 mM NaCl, pH 7,2, liquid or lyophilized, 0,01% NaN <sub>3</sub>  |
| Target-Kategorie       | Rat   |
| Antibody Type          | Polyclonal Antibody   |
| Application Verdünnung | WB: 1:2,000 - 1:10,000  |
| Anwendungsbeschreibung | Anti-Rat IgG antibody is suitable for use in ELISA, immunohistochemistry, and western blot. Specific conditions for reactivity should be optimized by the end user.   |