

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Goat IgM (μ)-unconj., MinX none, Polyclonal DNA-SEC-182522

| | |
|--------------------------|---|
| Artikelname | Rabbit IgG anti-Goat IgM (μ)-unconj., MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-182522 |
| Hersteller Artikelnummer | SEC-182522 |
| Alternativnummer | DNA-SEC-182522 |
| Hersteller | dianova |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | WB, IHC, ELISA |
| Spezies Reaktivität | Goat |
| Immunogen | Goat IgM mu heavy chain |
| Konjugation | Unconjugated |
| Produktbeschreibung | Anti-Goat IgM antibody specifically detects goat IgM. Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 ... |
| Klonalität | Polyclonal |
| Konzentration | 10.0 mg/mL |
| Isotyp | Ig |
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|------------------------|--|
| Reinheit | This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by |
| Formel | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target-Kategorie | Goat |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:2,000 - 1:10,000 |
| Anwendungsbeschreibung | Anti-Goat IgG IgM antibody is suitable for ELISA, western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency. |