

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit IgG anti-Guinea Pig IgG (H+L)-unconj., MinX none, Polyclonal DNA-SEC-182535

|                          |   |
|--------------------------|---|
| Artikelname              | Rabbit IgG anti-Guinea Pig IgG (H+L)-unconj., MinX none, Polyclonal   |
| Artikelnummer            | DNA-SEC-182535  |
| Hersteller Artikelnummer | SEC-182535  |
| Alternativnummer         | DNA-SEC-182535  |
| Hersteller               | dianova   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | WB, IHC, ELISA  |
| Spezies Reaktivität      | Guinea pig  |
| Immunogen                | Guinea Pig IgG whole molecule   |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | Anti-Guinea Pig IgG Antibody generated in rabbit detects guinea pig IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as we... |
| Klonalität               | Polyclonal  |
| Konzentration            | 10.0 mg/mL  |
| Isotyp                   | Ig  |
| Puffer                   | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                        |  |
|------------------------|--|
| Reinheit               | This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by |
| Formel                 | 20 mM K <sub>3</sub> PO <sub>4</sub> , 150 mM NaCl, pH 7,2, lyophilisate, Azide/BSA free   |
| Target-Kategorie       | Guinea Pig   |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | WB: 1:2,000 - 1:10,000   |
| Anwendungsbeschreibung | Anti-Guinea Pig IgG antibody is suitable for ELISA, western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency.  |