

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Horse IgG (H+L)-unconj., MinX none, Polyclonal DNA-SEC-182548

| | |
|--------------------------|---|
| Artikelname | Goat IgG anti-Horse IgG (H+L)-unconj., MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-182548 |
| Hersteller Artikelnummer | SEC-182548 |
| Alternativnummer | DNA-SEC-182548 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | WB, IHC, ELISA |
| Spezies Reaktivität | Equine |
| Immunogen | Horse IgG whole molecule |
| Konjugation | Unconjugated |
| Produktbeschreibung | Anti-Horse IgG Antibody generated in goat detects horse IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi ... |
| Klonalität | Polyclonal |
| Konzentration | 10.0 mg/mL |
| Isotyp | Ig |
| Puffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|------------------------|--|
| Reinheit | This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by |
| Formel | 10 mM NaPO ₄ , 150 mM NaCl, pH 7,2, lyophilisate, Azide/BSA free |
| Target-Kategorie | Horse |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:2,000 - 1:10,000 |
| Anwendungsbeschreibung | Anti-Horse IgG antibody is suitable for ELISA, western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency. |