

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat IgG anti-Human IgG (Fc)-unconj., MinX none, Polyclonal DNA-SEC-182562

|                          |   |
|--------------------------|---|
| Artikelname              | Goat IgG anti-Human IgG (Fc)-unconj., MinX none, Polyclonal   |
| Artikelnummer            | DNA-SEC-182562  |
| Hersteller Artikelnummer | SEC-182562  |
| Alternativnummer         | DNA-SEC-182562  |
| Hersteller               | dianova   |
| Wirt                     | Goat  |
| Kategorie                | Antikörper  |
| Applikation              | WB, IHC, ELISA  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human IgG F(c) fragment   |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | Anti-Human IgG F(c) generated in goat detects Human F(c). A proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of ... |
| Klonalität               | Polyclonal  |
| Konzentration            | 10.0 mg/mL  |
| Isotyp                   | Ig  |
| Puffer                   | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |

|                        |  |
|------------------------|--|
| Reinheit               | Anti-HUMAN IgG F(c) (GOAT) Antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer |
| Formel                 | 10 mM NaPO <sub>4</sub> , 150 mM NaCl, pH 7,2, lyophilisate, Azide/BSA free  |
| Target-Kategorie       | Human  |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | WB: 1:2,000 - 1:10,000   |
| Anwendungsbeschreibung | Anti-Human IgG F(c) fragment is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.            |