

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Human IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated DNA-SEC-182562

| | |
|--------------------------|---|
| Artikelname | Goat IgG anti-Human IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated |
| Artikelnummer | DNA-SEC-182562 |
| Hersteller Artikelnummer | SEC-182562 |
| Alternativnummer | DNA-SEC-182562 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | WB, IHC, ELISA |
| Spezies Reaktivität | Human |
| Immunogen | Human IgG F(c) fragment |
| Konjugation | Unconjugated |
| Produktbeschreibung | Anti-Human IgG F(c) generated in goat detects Human F(c). A proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of ... |
| Klonalität | Polyclonal |
| Konzentration | 10.0 mg/mL |
| Isotyp | Ig |

| | |
|------------------------|--|
| Puffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | Anti-HUMAN IgG F(c) (GOAT) Antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer |
| Formel | 10 mM NaPO ₄ , 150 mM NaCl, pH 7,2, lyophilisate, Azide/BSA free |
| Target-Kategorie | Human |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Anwendungsbeschreibung | Anti-Human IgG F(c) fragment is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user. |