

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat IgG anti-Donkey IgG (H+L)-unconj., MinX none, Polyclonal DNA-SEC-182637

|                          |   |
|--------------------------|---|
| Artikelname              | Goat IgG anti-Donkey IgG (H+L)-unconj., MinX none, Polyclonal   |
| Artikelnummer            | DNA-SEC-182637  |
| Hersteller Artikelnummer | SEC-182637  |
| Alternativnummer         | DNA-SEC-182637  |
| Hersteller               | dianova   |
| Wirt                     | Goat  |
| Kategorie                | Antikörper  |
| Applikation              | WB, IHC, ELISA  |
| Spezies Reaktivität      | Donkey  |
| Immunogen                | Donkey IgG whole molecule   |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | Anti-Donkey IgG Antibody generated in goat detects donkey IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fung... |
| Klonalität               | Polyclonal  |
| Konzentration            | 10.0 mg/mL  |
| Isotyp                   | Ig  |
| Puffer                   | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                        |  |
|------------------------|--|
| Reinheit               | Anti-DONKEY IgG (H&L) (GOAT) Antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buff |
| Formel                 | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3   |
| Target-Kategorie       | Donkey   |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | WB: 1:2,000 - 1:10,000   |
| Anwendungsbeschreibung | Anti-DONKEY IgG (H&L) (GOAT) Antibody is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency. |