

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit F(ab)2 anti-Goat IgG (H+L)-FITC, MinX none, Polyclonal DNA-SEC-182668

| | |
|--------------------------|---|
| Artikelname | Rabbit F(ab)2 anti-Goat IgG (H+L)-FITC, MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-182668 |
| Hersteller Artikelnummer | SEC-182668 |
| Alternativnummer | DNA-SEC-182668 |
| Hersteller | dianova |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | FC, IF, FLISA |
| Spezies Reaktivität | Goat |
| Immunogen | Goat IgG whole molecule |
| Konjugation | FITC |
| Produktbeschreibung | F(ab)2 Anti-Goat IgG Fluorescein Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunoche... |
| Klonalität | Polyclonal |
| Konzentration | 10.0 mg/mL |
| Isotyp | Ig |

| | |
|------------------------|---|
| Puffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | This product is a F(ab)2 fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis a |
| Formel | 10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target-Kategorie | Goat |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500-1:2,500, Fluorochrome Protein Value: 4.0, IF Microscopy Dilution: 1:1,000-1:5,000 |
| Anwendungsbeschreibung | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. |