

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Human IgG (H+L)-HRPO, MinX none, Polyclonal DNA-SEC-182689

| | |
|--------------------------|---|
| Artikelname | Goat F(ab)2 anti-Human IgG (H+L)-HRPO, MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-182689 |
| Hersteller Artikelnummer | SEC-182689 |
| Alternativnummer | DNA-SEC-182689 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | WB, IHC, ELISA |
| Spezies Reaktivität | Human |
| Immunogen | Human IgG whole molecule |
| Konjugation | HRP |
| Produktbeschreibung | F(ab)2 Anti-Human IgG (H&L) Peroxidase Antibody generated in goat detects immunoglobulin g from human, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum i... |
| Klonalität | Polyclonal |
| Konzentration | 10.0 mg/mL |
| Isotyp | Ig |
| Puffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|------------------------|---|
| Reinheit | This product is a F(ab)2 fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis a |
| Formel | 10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin |
| Target-Kategorie | Human |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:1,000 - 1:10,000 |
| Anwendungsbeschreibung | F(ab)2 Anti-Human IgG (H&L) Peroxidase secondary antibody reagents are ideal for ELISA, western blotting, Immunohistochemistry, Fluorescence Microscopy, Flow Cytometry as well as other antibody detection methods. |