

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat F(ab)2 anti-Mouse IgG (H+L)-unconj., MinX none, Polyclonal DNA-SEC-182693

|                          |   |
|--------------------------|---|
| Artikelname              | Goat F(ab)2 anti-Mouse IgG (H+L)-unconj., MinX none, Polyclonal   |
| Artikelnummer            | DNA-SEC-182693  |
| Hersteller Artikelnummer | SEC-182693  |
| Alternativnummer         | DNA-SEC-182693  |
| Hersteller               | dianova   |
| Wirt                     | Goat  |
| Kategorie                | Antikörper  |
| Applikation              | WB, IHC, ELISA  |
| Spezies Reaktivität      | Mouse   |
| Immunogen                | Mouse IgG whole molecule  |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | F(ab)2 Anti-Mouse IgG Antibody generated in goat was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain im... |
| Klonalität               | Polyclonal  |
| Konzentration            | 10.0 mg/mL  |
| Isotyp                   | Ig  |
| Puffer                   | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |

|                        |  |
|------------------------|--|
| Reinheit               | This product is a F(ab)2 fragment of IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by chromatographic separation |
| Formel                 | 10 mM NaPO4, 150 mM NaCl, pH 7.2, lyophilisate, Azide/BSA free   |
| Target-Kategorie       | Mouse  |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | WB: 1:2,000 - 1:10,000   |
| Anwendungsbeschreibung | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of Fc mediated binding, lot-to-lot consistency, high titer and specificity. The maxi     |