

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Chicken IgM (μ)-HRPO, MinX none, Polyclonal DNA-SEC-182780

| | |
|--------------------------|---|
| Artikelname | Goat IgG anti-Chicken IgM (μ)-HRPO, MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-182780 |
| Hersteller Artikelnummer | SEC-182780 |
| Alternativnummer | DNA-SEC-182780 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | ELISA, WB |
| Spezies Reaktivität | Gallus |
| Immunogen | Chicken IgM whole molecule |
| Konjugation | HRP |
| Produktbeschreibung | Anti-Chicken IgM antibody specifically detects chicken IgM. Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently link... |
| Klonalität | Polyclonal |
| Konzentration | 1.0 mg/mL |
| Isotyp | Ig |
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|------------------------|--|
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a sing |
| Formel | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin |
| Target-Kategorie | Chicken |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:1,000 - 1:10,000 |
| Anwendungsbeschreibung | Anti-Chicken IgM Peroxidase Conjugated antibody has been tested by ELISA and western blot. This product has been assayed against 1.0 ug of Chicken IgM in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code A |