

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Chicken IgG (H+L)-Alk. Phos., MinX Bo,Go,Gp,Hm,Ho,Hu,Ms,Rb,Rt,Sh, ALP, Polyclonal DNA-SEC-182784

| | |
|--------------------------|---|
| Artikelname | Goat IgG anti-Chicken IgG (H+L)-Alk. Phos., MinX Bo,Go,Gp,Hm,Ho,Hu,Ms,Rb,Rt,Sh, ALP, Polyclonal |
| Artikelnummer | DNA-SEC-182784 |
| Hersteller Artikelnummer | SEC-182784 |
| Alternativnummer | DNA-SEC-182784 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | DOT, ELISA, WB |
| Spezies Reaktivität | Gallus |
| Immunogen | Anti-Chicken IgG whole molecule was produced by repeated immunization with Chicken IgG whole molecule in goat. |
| Konjugation | ALP |
| Produktbeschreibung | Anti-Chicken IgG Alkaline Phosphatase Antibody generated in goat detects chicken IgY. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, ba... |
| Klonalität | Polyclonal |
| Konzentration | 1.0 mg/mL |

| | |
|------------------------|--|
| Isotyp | Ig |
| Puffer | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0 |
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by conjugation to fluorochrome and extensive dialysis against the buffer stated above. Assay by immunoelectrophores |
| Formel | 50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3 |
| Target-Kategorie | Chicken |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:500 - 1:2,500 |
| Anwendungsbeschreibung | Anti-Chicken IgG Alkaline Phosphatase Conjugate has been tested by ELISA, dot blot, and western blot. This product is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditio |