

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit IgG anti-Chicken IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated DNA-SEC-182789

|                          |   |
|--------------------------|---|
| Artikelname              | Rabbit IgG anti-Chicken IgG (Fc)-unconj., MinX none, Polyclonal ,<br>Unconjugated   |
| Artikelnummer            | DNA-SEC-182789  |
| Hersteller Artikelnummer | SEC-182789  |
| Alternativnummer         | DNA-SEC-182789  |
| Hersteller               | dianova   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | DOT, ELISA, WB  |
| Spezies Reaktivität      | Gallus  |
| Immunogen                | Chicken IgG F(c) fragment   |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | Anti-Chicken IgG F(c) generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of chicken IgG a... |
| Klonalität               | Polyclonal  |
| Konzentration            | 2.1 mg/mL   |
| Isotyp                   | Ig  |

|                        |  |
|------------------------|--|
| Puffer                 | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a sing |
| Formel                 | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3   |
| Target-Kategorie       | Chicken  |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1000-1:5000, Western Blot Dilution: 1:2000-1:10,000   |
| Anwendungsbeschreibung | Anti-Chicken IgG F(c) antibody has been tested by ELISA, dot blot, and western blot and is suitable for immunohistochemistry, as well as other assays requiring lot-to-lot consistency.  |