

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Dog IgG (Fc)-Alk. Phos., MinX none, ALP, Polyclonal DNA-SEC-182817

Artikelname	Goat IgG anti-Dog IgG (Fc)-Alk. Phos., MinX none, ALP, Polyclonal
Artikelnummer	DNA-SEC-182817
Hersteller Artikelnummer	SEC-182817
Alternativnummer	DNA-SEC-182817
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	WB, IHC, ELISA
Spezies Reaktivität	Canine
Immunogen	Dog IgG F(c) fragment
Konjugation	ALP
Produktbeschreibung	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglu...
Klonalität	Polyclonal
Konzentration	0.61 mg/mL
Isotyp	Ig

Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Dog IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single p
Formel	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target-Kategorie	Dog
Antibody Type	Polyclonal Antibody
Application Verdünnung	WB: 1:500 - 1:2,500
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Dog IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,000 to 1:5,000 of the reconstitution concent