

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Dog IgG (Fc)-Biotin, MinX none, Polyclonal DNA-SEC-182819

Artikelname	Goat IgG anti-Dog IgG (Fc)-Biotin, MinX none, Polyclonal
Artikelnummer	DNA-SEC-182819
Hersteller Artikelnummer	SEC-182819
Alternativnummer	DNA-SEC-182819
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA
Spezies Reaktivität	Canine
Immunogen	Dog IgG F(c) fragment
Konjugation	Biotin
Produktbeschreibung	Anti-DOG IgG F(c) (GOAT) Antibody generated in Goats detects specifically Dog IgG F(c). Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, ...
Klonalität	Polyclonal
Konzentration	2.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Reinheit	Anti-DOG IgG F(c) (GOAT) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Dog IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis r
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target-Kategorie	Dog
Antibody Type	Polyclonal Antibody
Application Verdünnung	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	Anti-DOG IgG F(c) Biotin Conjugated Antibody has been tested by ELISA and is assayed against 1.0 ug of Dog IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin S000-03 and ABTS (2,2-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid])