

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Goat IgG (H+L)-ATTO 550, MinX Ms,Rb, ATTO-550, Polyclonal DNA-SEC-182868

| | |
|--------------------------|---|
| Artikelname | Rabbit IgG anti-Goat IgG (H+L)-ATTO 550, MinX Ms,Rb, ATTO-550, Polyclonal |
| Artikelnummer | DNA-SEC-182868 |
| Hersteller Artikelnummer | SEC-182868 |
| Alternativnummer | DNA-SEC-182868 |
| Hersteller | dianova |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | WB |
| Spezies Reaktivität | Goat |
| Immunogen | Goat IgG whole molecule |
| Konjugation | ATTO-550 |
| Produktbeschreibung | Anti-Goat IgG ATTO dye Antibody generated in rabbit detects goat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well ... |
| Klonalität | Polyclonal |
| Konzentration | 1.0 mg/mL |
| Isotyp | Ig |

| | |
|------------------------|--|
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | Goat IgG (H&L) conjugated by ATTO 550 was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophore |
| Formel | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target-Kategorie | Goat |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | FLISA Dilution: >1:20,000, Fluorochrome Protein Value: 1.9, IF Microscopy Dilution: >1:5,000, Western Blot Dilution: >1:10,000 |
| Anwendungsbeschreibung | Anti-Goat IgG (H&L) conjugated by ATTO 550 has been tested by Western blot and is designed for STED microscopy, FRET, immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable f |