

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Donkey IgG anti-Goat IgG (H+L)-Alk. Phos., MinX Hm,Rb,Rt,Ck,Gp,Ho,Ms, ALP, Polyclonal , AP DNA-SEC-182888**

|                          |   |
|--------------------------|---|
| Artikelname              | Donkey IgG anti-Goat IgG (H+L)-Alk. Phos., MinX Hm,Rb,Rt,Ck,Gp,Ho,Ms, ALP, Polyclonal , AP  |
| Artikelnummer            | DNA-SEC-182888  |
| Hersteller Artikelnummer | SEC-182888  |
| Alternativnummer         | DNA-SEC-182888  |
| Hersteller               | dianova   |
| Wirt                     | Donkey  |
| Kategorie                | Antikörper  |
| Applikation              | ELISA   |
| Spezies Reaktivität      | Goat  |
| Immunogen                | Goat IgG whole molecule   |
| Konjugation              | AP  |
| Produktbeschreibung      | Anti-Goat IgG Alkaline Phosphatase Antibody generated in donkey detects goat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacter... |
| Klonalität               | Polyclonal  |
| Konzentration            | 1.0 mg/mL   |
| Isotyp                   | Ig  |

|                        |  |
|------------------------|--|
| Puffer                 | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0  |
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single  |
| Formel                 | 50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,1% NaN3   |
| Target-Kategorie       | Goat   |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | ELISA Dilution: 1:3,000 - 1:50,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,000   |
| Anwendungsbeschreibung | Anti-Goat IgG Alkaline Phosphatase Conjugated Antibody has been tested by ELISA and is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody-based enzymatic assays requiring |