

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Horse IgG (Fc)-Biotin, MinX none, Polyclonal DNA-SEC-182970

Artikelname	Goat IgG anti-Horse IgG (Fc)-Biotin, MinX none, Polyclonal
Artikelnummer	DNA-SEC-182970
Hersteller Artikelnummer	SEC-182970
Alternativnummer	DNA-SEC-182970
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	WB, IHC, ELISA
Spezies Reaktivität	Equine
Immunogen	Horse IgG F(c) fragment
Konjugation	Biotin
Produktbeschreibung	Anti-Horse IgG F(c) generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of horse IgG and oft...
Klonalität	Polyclonal
Konzentration	2.0 mg/mL
Isotyp	Ig
Puffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single
Formel	10 mM NaPO ₄ ,150 mM NaCl,pH 7,2,lyophilisate,0,01% Thimerosal
Target-Kategorie	Horse
Antibody Type	Polyclonal Antibody
Application Verdünnung	WB: 1:2,000 - 1:10,000
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Horse IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin S000-03 and ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 minutes at room