

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit IgG anti-Horse IgG (Fc)-HRPO, MinX none, Polyclonal DNA-SEC-182979

|                          |   |
|--------------------------|---|
| Artikelname              | Rabbit IgG anti-Horse IgG (Fc)-HRPO, MinX none, Polyclonal  |
| Artikelnummer            | DNA-SEC-182979  |
| Hersteller Artikelnummer | SEC-182979  |
| Alternativnummer         | DNA-SEC-182979  |
| Hersteller               | dianova   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | WB, IHC, ELISA  |
| Spezies Reaktivität      | Equine  |
| Immunogen                | Horse IgG F(c) fragment   |
| Konjugation              | HRP   |
| Produktbeschreibung      | Anti-Horse IgG F(c) generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of horse IgG and o... |
| Klonalität               | Polyclonal  |
| Konzentration            | 1.5 mg/mL   |
| Isotyp                   | Ig  |
| Puffer                   | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                        |  |
|------------------------|--|
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single |
| Formel                 | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin   |
| Target-Kategorie       | Horse  |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | WB: 1:1,000 - 1:10,000   |
| Anwendungsbeschreibung | This product has been assayed against 1.0 ug of Horse IgG in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to     |