

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat IgG anti-Human IgG (H+L)-Alk. Phos., MinX Ms, ALP, Polyclonal , AP DNA-SEC-183039

|                          |   |
|--------------------------|---|
| Artikelname              | Goat IgG anti-Human IgG (H+L)-Alk. Phos., MinX Ms, ALP, Polyclonal , AP   |
| Artikelnummer            | DNA-SEC-183039  |
| Hersteller Artikelnummer | SEC-183039  |
| Alternativnummer         | DNA-SEC-183039  |
| Hersteller               | dianova   |
| Wirt                     | Goat  |
| Kategorie                | Antikörper  |
| Applikation              | DOT, ELISA  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human IgG whole molecule  |
| Konjugation              | AP  |
| Produktbeschreibung      | Anti-Human IgG (H&L) Alkaline Phosphatase generated in goat detects human Immunoglobulin G (IgG), both heavy and light chains of the antibody molecule are present. It is a protein complex composed of four peptide chains - two identical heavy chains a... |
| Klonalität               | Polyclonal  |
| Konzentration            | 1.0 mg/mL   |
| Isotyp                   | Ig  |

|                        |  |
|------------------------|--|
| Puffer                 | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0  |
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single |
| Formel                 | 50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3  |
| Target-Kategorie       | Human  |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500   |
| Anwendungsbeschreibung | Anti-Human IgG (H&L) Alkaline Phosphatase has been tested by dot blot and ELISA and is assayed against 1.0 ug of Human IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code NPP-10 as a substrate for 30 minutes at room temperature. A w   |