

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Rabbit IgG anti-Human IgG (Fc)-Alk. Phos., MinX none, ALP, Polyclonal , AP DNA-SEC-183079**

|                          |   |
|--------------------------|---|
| Artikelname              | Rabbit IgG anti-Human IgG (Fc)-Alk. Phos., MinX none, ALP, Polyclonal , AP  |
| Artikelnummer            | DNA-SEC-183079  |
| Hersteller Artikelnummer | SEC-183079  |
| Alternativnummer         | DNA-SEC-183079  |
| Hersteller               | dianova   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | DOT, ELISA  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Anti-Human IgG F(c) was produced by repeated immunization with Human Fc fragment in rabbit.   |
| Konjugation              | AP  |
| Produktbeschreibung      | Anti-Human IgG F(c) Alkaline Phosphatase Conjugated generated in rabbit detects Human F(c). A proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH.... |
| Klonalität               | Polyclonal  |
| Konzentration            | 1.0 mg/mL   |

|                        |  |
|------------------------|--|
| Isotyp                 | Ig   |
| Puffer                 | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0  |
| Reinheit               | Anti-Human IgG F(c) was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a |
| Formel                 | 50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0.05% NaN3  |
| Target-Kategorie       | Human  |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500   |
| Anwendungsbeschreibung | Anti-Human IgG F(c) Alkaline Phosphatase Conjugated antibody has been tested by dot blot and ELISA and is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for react |