

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Mouse IgM (μ)-FITC, MinX none, Polyclonal DNA-SEC-183168

| | |
|--------------------------|---|
| Artikelname | Goat IgG anti-Mouse IgM (μ)-FITC, MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-183168 |
| Hersteller Artikelnummer | SEC-183168 |
| Alternativnummer | DNA-SEC-183168 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | WB |
| Spezies Reaktivität | Mouse |
| Immunogen | Mouse IgM whole molecule |
| Konjugation | FITC |
| Produktbeschreibung | Anti-Mouse IgM antibody specifically detects mouse IgM. Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking ... |
| Klonalität | Polyclonal |
| Konzentration | 2.0 mg/mL |
| Isotyp | Ig |
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|------------------------|---|
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single |
| Formel | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target-Kategorie | Mouse |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 3.7, IF Microscopy Dilution: 1:1,000 - 1:5,000 |
| Anwendungsbeschreibung | Anti-Mouse IgM antibody has been tested by western blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor |