

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Rabbit IgG (F(ab)2)-HRPO, MinX none, Polyclonal DNA-SEC-183366

| | |
|--------------------------|---|
| Artikelname | Goat IgG anti-Rabbit IgG (F(ab)2)-HRPO, MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-183366 |
| Hersteller Artikelnummer | SEC-183366 |
| Alternativnummer | DNA-SEC-183366 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | ELISA |
| Spezies Reaktivität | Rabbit |
| Immunogen | Rabbit IgG F(ab)2 fragment |
| Konjugation | HRP |
| Produktbeschreibung | Anti-Rabbit IgG F(ab)2 Antibody generated in goat recognizes the dimeric Fab portion of the rabbit IgG molecule. Rabbit IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controll... |
| Klonalität | Polyclonal |
| Konzentration | 2.0 mg/mL |
| Isotyp | Ig |
| Puffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|------------------------|--|
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a singl |
| Formel | 10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin |
| Target-Kategorie | Rabbit |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000 |
| Anwendungsbeschreibung | Anti-Rabbit IgG F(ab)2 peroxidase conjugated antibody has been tested by ELISA and is assayed against 1.0 ug of Rabbit IgG in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 3 |