

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Donkey IgG anti-Rabbit IgG (Fc)-unconj., MinX none, Polyclonal DNA-SEC-183420

| | |
|--------------------------|---|
| Artikelname | Donkey IgG anti-Rabbit IgG (Fc)-unconj., MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-183420 |
| Hersteller Artikelnummer | SEC-183420 |
| Alternativnummer | DNA-SEC-183420 |
| Hersteller | dianova |
| Wirt | Donkey |
| Kategorie | Antikörper |
| Applikation | WB |
| Spezies Reaktivität | Rabbit |
| Immunogen | Rabbit IgG F(c) fragment |
| Konjugation | Unconjugated |
| Produktbeschreibung | Anti-Rabbit IgG F(c) generated in donkey is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of rabbit IgG and... |
| Klonalität | Polyclonal |
| Konzentration | 2 mg/mL |
| Isotyp | Ig |
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|------------------------|--|
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Rabbit IgG, Rabbit IgG F(c) and Rabb |
| Formel | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3 |
| Target-Kategorie | Rabbit |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:2,000 - 1:20,000 |
| Anwendungsbeschreibung | Anti-Rabbit IgG F(c) antibody is tested by western blot and suitable for ELISA and immunohistochemistry, as well as other assays requiring lot-to-lot consistency. |