

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Rabbit IgG anti-Rat IgG (F(ab)2)-unconj., MinX none, Polyclonal , Unconjugated DNA-SEC-183493**

|                          |   |
|--------------------------|---|
| Artikelname              | Rabbit IgG anti-Rat IgG (F(ab)2)-unconj., MinX none, Polyclonal ,<br>Unconjugated   |
| Artikelnummer            | DNA-SEC-183493  |
| Hersteller Artikelnummer | SEC-183493  |
| Alternativnummer         | DNA-SEC-183493  |
| Hersteller               | dianova   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, WB   |
| Spezies Reaktivität      | Rat   |
| Immunogen                | Rat IgG F(ab)2 fragment   |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | Anti-Rat IgG F(ab)2 Antibody generated in rabbit recognizes the dimeric Fab portion of the rat IgG molecule. Rat IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled cond... |
| Klonalität               | Polyclonal  |
| Konzentration            | 7.3 mg/mL   |
| Isotyp                   | Ig  |

|                        |  |
|------------------------|--|
| Puffer                 | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single p |
| Formel                 | 10 mM NaPO <sub>4</sub> , 150 mM NaCl, pH 7,2, sterile filtered, 0,01% NaN <sub>3</sub>  |
| Target-Kategorie       | Rat  |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000  |
| Anwendungsbeschreibung | Anti-Rat IgG F(ab) <sub>2</sub> antibody has been tested by ELISA and western blot and is suitable for use in immunohistochemistry. Specific conditions for reactivity should be optimized by the end user.  |