

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Hamster armenian IgG (H+L)-Biotin, MinX none, Polyclonal DNA-SEC-183647

| | |
|--------------------------|---|
| Artikelname | Goat IgG anti-Hamster armenian IgG (H+L)-Biotin, MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-183647 |
| Hersteller Artikelnummer | SEC-183647 |
| Alternativnummer | DNA-SEC-183647 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | ELISA, WB |
| Spezies Reaktivität | Hamster |
| Immunogen | Armenian Hamster IgG whole molecule |
| Konjugation | Biotin |
| Produktbeschreibung | Anti-Armenian Hamster IgG Biotin Antibody generated in goat detects Armenian Hamster IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses... |
| Klonalität | Polyclonal |
| Konzentration | 1.0 mg/mL |
| Isotyp | Ig |

| | |
|------------------------|--|
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Armenian Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted |
| Formel | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target-Kategorie | Armenian Hamster |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:2,000 - 1:10,000 |
| Anwendungsbeschreibung | Biotin Anti-Hamster IgG Secondary Antibody has been tested by ELISA and western blot and is designed for Western Blotting, ELISA and Immunohistochemistry. Biotin conjugated secondary antibodies can also be used for a variety of other applications such as |