

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Donkey F(ab)2 anti-Goat IgG (H+L)-RPE, MinX Ck,Gp,Hm,Ho,Hu,Ms,Rb,Rt, Polyclonal DNA-SEC-183694**

|                          |   |
|--------------------------|---|
| Artikelname              | Donkey F(ab)2 anti-Goat IgG (H+L)-RPE, MinX Ck,Gp,Hm,Ho,Hu,Ms,Rb,Rt, Polyclonal   |
| Artikelnummer            | DNA-SEC-183694  |
| Hersteller Artikelnummer | SEC-183694  |
| Alternativnummer         | DNA-SEC-183694  |
| Hersteller               | dianova   |
| Wirt                     | Donkey  |
| Kategorie                | Antikörper  |
| Applikation              | DOT, WB   |
| Spezies Reaktivität      | Goat  |
| Immunogen                | Anti-Goat IgG was produced by repeated immunization with goat IgG whole molecule in donkey.   |
| Konjugation              | RPE   |
| Produktbeschreibung      | F(ab)2 Anti-Goat IgG Phycoerythrin Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunoc... |
| Klonalität               | Polyclonal  |
| Konzentration            | 1.032 mg/ml   |

|                        |  |
|------------------------|--|
| Isotyp                 | Ig   |
| Puffer                 | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Ass |
| Formel                 | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3   |
| Target-Kategorie       | Goat   |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | WB: User Optimized   |
| Anwendungsbeschreibung | F(ab)2 Anti-Goat IgG Phycoerythrin Antibody has been tested by dot blot and western blot and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels |