

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat F(ab)2 anti-Guinea Pig IgG (H+L)-unconj., MinX none, Polyclonal DNA-SEC-183695

|                          |   |
|--------------------------|---|
| Artikelname              | Goat F(ab)2 anti-Guinea Pig IgG (H+L)-unconj., MinX none, Polyclonal  |
| Artikelnummer            | DNA-SEC-183695  |
| Hersteller Artikelnummer | SEC-183695  |
| Alternativnummer         | DNA-SEC-183695  |
| Hersteller               | dianova   |
| Wirt                     | Goat  |
| Kategorie                | Antikörper  |
| Applikation              | WB, IHC, ELISA  |
| Spezies Reaktivität      | Guinea pig  |
| Immunogen                | Guinea Pig IgG whole molecule   |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | F(ab)2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and exper... |
| Klonalität               | Polyclonal  |
| Konzentration            | 1.0 mg/mL   |
| Isotyp                   | Ig  |

|                        |  |
|------------------------|--|
| Puffer                 | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipit |
| Formel                 | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3   |
| Target-Kategorie       | Guinea Pig   |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | WB: 1:2,000 - 1:10,000   |
| Anwendungsbeschreibung | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maxi |