

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit F(ab)2 anti-Horse IgG (H+L)-Biotin, MinX none, Polyclonal DNA-SEC-183716

| | |
|--------------------------|---|
| Artikelname | Rabbit F(ab)2 anti-Horse IgG (H+L)-Biotin, MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-183716 |
| Hersteller Artikelnummer | SEC-183716 |
| Alternativnummer | DNA-SEC-183716 |
| Hersteller | dianova |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | WB, IHC, ELISA |
| Spezies Reaktivität | Equine |
| Immunogen | Horse IgG whole molecule |
| Konjugation | Biotin |
| Produktbeschreibung | F(ab)2 Anti-Horse IgG Biotin Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemica... |
| Klonalität | Polyclonal |
| Konzentration | 1.0 mg/mL |
| Isotyp | Ig |
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|------------------------|--|
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. As |
| Formel | 20 mM K ₃ PO ₄ , 150 mM NaCl, pH 7,2, lyophilisate, 0,01% NaN ₃ |
| Target-Kategorie | Horse |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:10,000 - 1:25,000 |
| Anwendungsbeschreibung | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and |