

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Human Lambda light chain-unconj., MinX none, Polyclonal DNA-SEC-183725

| | |
|--------------------------|---|
| Artikelname | Goat F(ab)2 anti-Human Lambda light chain-unconj., MinX none, Polyclonal |
| Artikelnummer | DNA-SEC-183725 |
| Hersteller Artikelnummer | SEC-183725 |
| Alternativnummer | DNA-SEC-183725 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | WB, IHC, ELISA |
| Spezies Reaktivität | Human |
| Immunogen | Human lambda light chain |
| Konjugation | Unconjugated |
| Produktbeschreibung | F(ab)2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and exper... |
| Klonalität | Polyclonal |
| Konzentration | 1.0 mg/mL |

| | |
|------------------------|--|
| Isotyp | Ig |
| Puffer | 0.125 M Sodium Borate, 0.075 M Sodium Chloride, 0.005 M EDTA, pH 8.0 |
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Ass |
| Formel | 125 mM Sodium Borate, 75 mM NaCl, 5 mM EDTA, pH 8,0, sterile filtered, 0,01% NaN ₃ |
| Target-Kategorie | Human |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:1,000 - 1:5,000 |
| Anwendungsbeschreibung | Suitable for highly specific immunological methods requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. |