

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat F(ab)2 anti-Human IgG (H)-unconj., MinX none, Polyclonal DNA-SEC-183726

|                          |   |
|--------------------------|---|
| Artikelname              | Goat F(ab)2 anti-Human IgG (H)-unconj., MinX none, Polyclonal   |
| Artikelnummer            | DNA-SEC-183726  |
| Hersteller Artikelnummer | SEC-183726  |
| Alternativnummer         | DNA-SEC-183726  |
| Hersteller               | dianova   |
| Wirt                     | Goat  |
| Kategorie                | Antikörper  |
| Applikation              | WB, IHC, ELISA  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human IgG gamma heavy chain   |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | F(ab)2 Anti-Human IgG (gamma chain) Antibody generated in goat detects human immunoglobulin g (gamma chain). Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins in humans, IgG is the most abundant antibody ... |
| Klonalität               | Polyclonal  |
| Konzentration            | 1.0 mg/mL   |
| Isotyp                   | Ig  |

|                        |  |
|------------------------|--|
| Puffer                 | 0.125 M Sodium Borate, 0.075 M Sodium Chloride, 0.005 M EDTA, pH 8.0   |
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Ass |
| Formel                 | 125 mM Sodium Borate, 75 mM NaCl, 5 mM EDTA, pH 8,0, sterile filtered, 0,01% NaN <sub>3</sub>  |
| Target-Kategorie       | Human  |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | WB: 1:2,000 - 1:10,000   |
| Anwendungsbeschreibung | Suitable for highly specific immunological methods requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.  |