

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Mouse IgG+IgM+IgA (H+L)-unconj., MinX Hu, Polyclonal DNA-SEC-183770

| | |
|--------------------------|---|
| Artikelname | Goat F(ab)2 anti-Mouse IgG+IgM+IgA (H+L)-unconj., MinX Hu, Polyclonal |
| Artikelnummer | DNA-SEC-183770 |
| Hersteller Artikelnummer | SEC-183770 |
| Alternativnummer | DNA-SEC-183770 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | ELISA |
| Spezies Reaktivität | Mouse |
| Immunogen | Mouse IgG IgA and IgM whole molecule |
| Konjugation | Unconjugated |
| Produktbeschreibung | F(ab)2 Anti-Mouse IgG IgA IgM (H&L) Antibody generated in goat detects reactivity to Mouse IgG, Mouse IgA, and Mouse IgM subclasses and chains. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary ... |
| Klonalität | Polyclonal |
| Konzentration | 1.0 mg/mL |
| Isotyp | Ig |

| | |
|------------------------|--|
| Puffer | 0.125 M Sodium Borate, 0.075 M Sodium Chloride, 0.005 M EDTA, pH 8.0 |
| Reinheit | This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single |
| Formel | 125 mM Sodium Borate, 75 mM NaCl, 5 mM EDTA, pH 8,0, sterile filtered, 0,01% Na ₃ N |
| Target-Kategorie | Mouse |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:2000-1:10,000 |
| Anwendungsbeschreibung | F(ab) ₂ Anti-Mouse IgG IgA IgM (H&L) Antibody has been tested by ELISA and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) m |