

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Rabbit IgG (F(ab)2)-Alk. Phos., MinX Bo, Ho, Hu, Ms, Rt, Sh, ALP, Polyclonal DNA-SEC-183837

| | |
|--------------------------|---|
| Artikelname | Goat F(ab)2 anti-Rabbit IgG (F(ab)2)-Alk. Phos., MinX Bo, Ho, Hu, Ms, Rt, Sh, ALP, Polyclonal |
| Artikelnummer | DNA-SEC-183837 |
| Hersteller Artikelnummer | SEC-183837 |
| Alternativnummer | DNA-SEC-183837 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | ELISA |
| Spezies Reaktivität | Rabbit |
| Immunogen | Rabbit IgG F(ab)2 fragment |
| Konjugation | ALP |
| Produktbeschreibung | F(ab)2 Anti-Rabbit IgG F(ab)2 Alkaline Phosphatase Antibody generated in goat detects Rabbit F(ab)2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthe... |
| Klonalität | Polyclonal |
| Konzentration | 0.5 mg/mL |

| | |
|------------------------|--|
| Isotyp | Ig |
| Puffer | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0 |
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. A |
| Formel | 50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3 |
| Target-Kategorie | Rabbit |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | WB: 1:500 - 1:2,500 |
| Anwendungsbeschreibung | F(ab)2 Anti-Rabbit IgG F(ab)2 Alkaline Phosphatase Antibody has been tested by ELISA and is assayed against 1.0 µg of Rabbit IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code NPP-10 as a substrate for 30 minutes at room temperatur |