

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Donkey F(ab)2 anti-Rabbit IgG (H+L)-unconj., MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh, Polyclonal DNA-SEC-183842**

|                          |   |
|--------------------------|---|
| Artikelname              | Donkey F(ab)2 anti-Rabbit IgG (H+L)-unconj., MinX<br>Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh, Polyclonal  |
| Artikelnummer            | DNA-SEC-183842  |
| Hersteller Artikelnummer | SEC-183842  |
| Alternativnummer         | DNA-SEC-183842  |
| Hersteller               | dianova   |
| Wirt                     | Donkey  |
| Kategorie                | Antikörper  |
| Applikation              | ELISA   |
| Spezies Reaktivität      | Rabbit  |
| Immunogen                | Rabbit IgG whole molecule   |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | F(ab)2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and exper... |
| Klonalität               | Polyclonal  |
| Konzentration            | 1.0 mg/mL   |

|                        |  |
|------------------------|--|
| Isotyp                 | Ig   |
| Puffer                 | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. A |
| Formel                 | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3   |
| Target-Kategorie       | Rabbit   |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | WB: 1:2,000-1:10,000   |
| Anwendungsbeschreibung | F(ab)2 Anti-Rabbit IgG Antibody has been tested by ELISA and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated bindi |